

IN THE CLAIMS:

The status of each claim that has been introduced in the above-referenced application is identified in the ensuing listing of the claims. This listing of the claims replaces all previously submitted claims listings.

1. (Currently amended) A method for performing an assay, comprising:  
applying a sample to an assay device carrying a plurality of reactive elements for assaying a plurality of analytes in the sample;  
concurrently evaluating the presence of ~~a the~~ plurality of analytes ~~in a single portion of a sample applied to a single assay device, at least one analyte~~ a combination of analytes of the plurality of analytes having known parameters that, in combination, are indicative of an acute metabolic or disease state; ~~substantially simultaneously~~ including concurrently determining concentrations rates of reaction of each type of reactive element of the plurality of reactive elements with its corresponding analyte of each of the plurality of analytes in the sample;  
continuing the ~~substantially simultaneous concurrent~~ determination until a combination of reaction rates provides a reliable indication ~~the at least one analyte has been reliably determined to be present in an amount indicative of a presence or absence of the acute~~ metabolic or disease state; and  
reporting ~~said reliable results of the concurrent determination of the presence of the plurality of analytes in an amount indicative of the metabolic or disease state.~~

2. (Currently amended) The method according to claim 1, wherein concurrently determining ~~evaluating the presence of at least one other analyte in the sample continues after the report of the reliable determination of an amount indicative of the acute metabolic or disease state in order to accurately determine the presence or concentration of the at least one other analyte~~ reporting results and further comprising reporting further results of the concurrent determination.

3. (Original) The method according to claim 1, comprising evaluating binding of the plurality of analytes to corresponding reactive elements over a plurality of time points.
4. (Canceled)
5. (Canceled)
6. (Currently amended) The method according to claim ~~5~~1, wherein each reactive element is substantially immobilized on a single waveguide surface.
7. (Canceled)
8. (Currently amended) The method according to ~~claim 4~~, claim 1, wherein the reactive elements are arranged in one or more patterns on a waveguide surface.
9. (Currently amended) The method according to ~~claim 4~~, claim 6, wherein the ~~substantially simultaneous concurrent~~ determination includes introducing a light beam including at least one wavelength appropriate for stimulating a light signal from the corresponding reactive element when the corresponding reactive element has coupled with the at least one analyte.
10. (Original) The method according to claim 9, wherein the light signal is indicative of a rate of reaction between the analyte of interest and the corresponding reactive element.
11. (Original) The method according to claim 10, wherein the substantially simultaneous determination includes measuring the light signal generated from the reaction of the at least one analyte with the corresponding reactive element.
12. (Canceled)

13. (Original) The method according to claim 1, wherein the at least one analyte is a marker released from cardiac tissue only after a myocardial infarction.

14. (Original) The method according to claim 13, wherein the marker comprises myoglobin.

15. (Original) The method according to claim 1, wherein the at least one analyte is a cardiac specific marker.

16. (Original) The method according to claim 15, wherein the at least one analyte comprises troponin.

17. (Original) The method according to claim 16, wherein the troponin comprises individual troponin subunits.

18. (Original) The method according to claim 16, wherein the troponin comprises a complex including at least one troponin subunit.

19. (Original) The method according to claim 16, wherein the troponin comprises at least one of native troponin and a modified troponin.

20. (Original) The method according to claim 15, wherein the at least one analyte comprises creatine kinase.

21. (Original) The method according to claim 20, wherein the creatine kinase comprises CK-MB.

22-35 (Canceled)